

# First-in-class Interleukin-15 Fusion Protein for the Treatment of Solid Tumors

## Overview

Drug Name	BioLink001
Description	BioLink001 is a tumor-targeting interleukin-15 (IL-15) fusion protein and is in early
	clinical development for the treatment of locally advanced/metastatic or
	relapsed/refractory solid tumors. BioLink001 was granted orphan drug designation
	by the FDA.
Target	Interleukin 15
Drug Modality	Fusion protein
Indication	Solid tumor
Product Category	Cancer immunotherapy
Mechanism of Action	Interleukin 15 replacements
Status	Phase I
Patent	Granted

## **Collaboration Opportunity**

Protheragen Inc. is actively seeking partnership for BioLink001. Potential collaboration can be strategic alliance, licensing, or marketing agreement. We look forward to hearing from you.

## Target

## **Interleukin 15**

Introduction

IL-15 is a cytokine expressed by monocytes, macrophages, dendritic cells (DC), keratinocytes, fibroblasts, and nerve cells. It binds to and signals through a



complex composed of the IL-2/IL-15 receptor beta chain (CD122) and the common         gamma chain (CD132). IL-15 effects on many cells of the immune system and can         promote the proliferation of B cells and the secretion of immunoglobulins. It is         essential for the ontogeny of natural killer (NK) cells and CD8+ T cells, inducing the         activation, proliferation, cytolytic activity and the production of cytokines such as         interferon-y (IFN-y). IL-15 has no significant effects on regulatory T cells (Tregs),         but plays an important role in supporting the long-term maintenance and high-         affinity response of T cells to malignant cells by acting as a survival factor for         memory CD8+ T cells.         In addition, IL-15 has pro-inflammatory potential. It is implicated in the         pathogenesis of several autoimmune diseases, such as rheumatoid arthritis,         psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted.         However, viral infection can cause its secretion.         Approved Name       Interleukin 15         Official Symbol       IL15         Gene Type       Protein coding         Synonyms       IL-15; MGC9721         Ensembl       ENSG0000164136         Gene ID       3600         mRNA Refseq       NM_000585; NM_172175         Protein Refseq       NP_000576; NP_751915         OMIM <td< th=""><th></th><th></th></td<>		
promote the proliferation of B cells and the secretion of immunoglobulins. It is essential for the ontogeny of natural killer (NK) cells and CD8+ T cells, inducing the activation, proliferation, cytolytic activity and the production of cytokines such as interferon-γ (IFN-γ). IL-15 has no significant effects on regulatory T cells (Tregs), but plays an important role in supporting the long-term maintenance and high- affinity response of T cells to malignant cells by acting as a survival factor for memory CD8+ T cells. In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL-15, MGC9721EnsemblENSG0000164136 Gene ID 3600 mRNA RefseqMN 000585; NM 172175Protein RefseqProtein RefseqNP 000576; NP 751915 OMIM 600554 UniProt IDQuestionP40933		complex composed of the IL-2/IL-15 receptor beta chain (CD122) and the common
Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG0000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		
activation, proliferation, cytolytic activity and the production of cytokines such as interferon-γ (IFN-γ). IL-15 has no significant effects on regulatory T cells (Tregs), but plays an important role in supporting the long-term maintenance and high- affinity response of T cells to malignant cells by acting as a survival factor for memory CD8+ T cells. In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG00000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		
<ul> <li>interferon-y (IFN-y). IL-15 has no significant effects on regulatory T cells (Tregs), but plays an important role in supporting the long-term maintenance and high- affinity response of T cells to malignant cells by acting as a survival factor for memory CD8+ T cells. In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.</li> <li>Approved Name Interleukin 15</li> <li>Official Symbol IL15</li> <li>Gene Type Protein coding</li> <li>Synonyms IL-15; MGC9721</li> <li>Ensembl ENSG0000164136</li> <li>Gene ID 3600</li> <li>mRNA Refseq NM_000585; NM_172175</li> <li>Protein Refseq NP_000576; NP_751915</li> <li>OMIM 600554</li> <li>UniProt ID P40933</li> </ul>		
but plays an important role in supporting the long-term maintenance and high- affinity response of T cells to malignant cells by acting as a survival factor for memory CD8+ T cells. In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG0000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		activation, proliferation, cytolytic activity and the production of cytokines such as
affinity response of T cells to malignant cells by acting as a survival factor for memory CD8+ T cells. In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, 		interferon- $\gamma$ (IFN- $\gamma$ ). IL-15 has no significant effects on regulatory T cells (Tregs),
memory CD8+ T cells.In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG0000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		but plays an important role in supporting the long-term maintenance and high-
In addition, IL-15 has pro-inflammatory potential. It is implicated in the pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG0000164136Gene ID3600mRNA RefseqNM_000585; NM_172175Protein RefseqNP_000576; NP_751915OMIM600554UniProt IDP40933		affinity response of T cells to malignant cells by acting as a survival factor for
pathogenesis of several autoimmune diseases, such as rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG00000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		memory CD8+ T cells.
psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted. However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG00000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		In addition, IL-15 has pro-inflammatory potential. It is implicated in the
However, viral infection can cause its secretion.Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG00000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		pathogenesis of several autoimmune diseases, such as rheumatoid arthritis,
Approved NameInterleukin 15Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG00000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		psoriasis, and pulmonary inflammatory diseases. IL-15 is normally not secreted.
Official SymbolIL15Gene TypeProtein codingSynonymsIL-15; MGC9721EnsemblENSG00000164136Gene ID3600mRNA RefseqNM 000585; NM 172175Protein RefseqNP 000576; NP 751915OMIM600554UniProt IDP40933		However, viral infection can cause its secretion.
Gene Type         Protein coding           Synonyms         IL-15; MGC9721           Ensembl         ENSG00000164136           Gene ID         3600           mRNA Refseq         NM_000585; NM_172175           Protein Refseq         NP_000576; NP_751915           OMIM         600554           UniProt ID         P40933	Approved Name	Interleukin 15
Synonyms         IL-15; MGC9721           Ensembl         ENSG00000164136           Gene ID         3600           mRNA Refseq         NM 000585; NM 172175           Protein Refseq         NP 000576; NP 751915           OMIM         600554           UniProt ID         P40933	Official Symbol	IL15
Ensembl         ENSG0000164136           Gene ID         3600           mRNA Refseq         NM 000585; NM 172175           Protein Refseq         NP 000576; NP 751915           OMIM         600554           UniProt ID         P40933	Gene Type	Protein coding
Gene ID         3600           mRNA Refseq         NM 000585; NM 172175           Protein Refseq         NP 000576; NP 751915           OMIM         600554           UniProt ID         P40933	Synonyms	IL-15; MGC9721
mRNA Refseq         NM_000585; NM_172175           Protein Refseq         NP_000576; NP_751915           OMIM         600554           UniProt ID         P40933	Ensembl	ENSG0000164136
Protein Refseq         NP 000576; NP 751915           OMIM         600554           UniProt ID         P40933	Gene ID	3600
OMIM         600554           UniProt ID         P40933	mRNA Refseq	<u>NM_000585; NM_172175</u>
UniProt ID         P40933	Protein Refseq	<u>NP_000576; NP_751915</u>
	OMIM	600554
Chromosome Location 4q31.21	UniProt ID	P40933
	Chromosome Location	4q31.21

# **Drug Modality**

## **Fusion Protein**

BioLink001 is an IL-15 fusion protein developed with global intellectual property rights. It is composed of an integrin-binding arginylglycylaspartic acid (RGD)-4C motif, linked with a human IgG1 Fc, and a modified sushi domain of human IL-15R $\alpha$  unit and a human IL-15. The RGD-4C motif of BioLink001 binds the  $\alpha\nu\beta3$ ,  $\alpha\nu\beta5$ , and  $\alpha\nu\beta6$  integrins, which are commonly overexpressed in solid tumors, leading to the tumor-targeting.

# Indication

#### Solid Tumors

Solid tumors are abnormal masses of tissue that usually do not contain cysts or areas of fluid. Solid tumors may be benign or malignant. Different types of solid tumors are named for the cell type that forms them, such as sarcomas, carcinomas, and lymphomas. Solid tumors account for approximately 90% of cancers in adults and make up about 30% of all cancers in children. There were an estimated 18.1 million cancer cases worldwide in 2020. The most common cancers are breast, lung, colon, rectum, and prostate cancers. The most common type of solid tumor found in children is the brain tumor.

Cancer treatment usually includes surgery, radiotherapy, chemotherapy, hormonal treatments, and targeted biological therapies. Cancer treatment decisions should take into account the stage and biology of the tumor, risk and prognosis of planned therapy, and the economic costs associated with therapy. The global solid tumor cancer treatment market was estimated at \$121.3 billion in 2018 and is projected to reach \$424.6 billion by 2027, increasing to CAGR by 15.0% from 2019 to 2027.

## **Mechanism of Action**

#### **Interleukin 15 Replacements**

IL-15 is essential in the growth, mobilization, and activation of immune cells such as NK cells, CD8+ T cells,  $\gamma\delta$  T cells and natural killer T (NKT) cells. IL-15 can also enhance the anti-tumor activity of immune cells, acts as a survival factor for CD8 memory T cells, and contributes to the long-term maintenance of high affinity T cell response to malignant cells. IL-15 is a promising immunotherapy agent due to its potent immune actions. BioLink001 is a tumor-targeting IL-15 fusion protein that is composed of an RGD-4C motif, a human IgG1 Fc, and a modified sushi domain of human IL-15R $\alpha$  unit and a human IL-15. BioLink001 targets cancer cells that overexpress integrins such as  $\alpha\nu\beta3$ ,  $\alpha\nu\beta5$ , and  $\alpha\nu\beta6$  by the integrin-binding RGD-4C motif and stimulates anticancer immunity by its IL-15 motif. This design could enhance anticancer effects and reduce systemic toxicity



of BioLink001 by localizing the role of IL-15 in the tumor microenvironment.

## Status

## The Status of BioLink001

In the ongoing phase 1 clinical trial in adult patients with solid tumors, BioLink001 as a single agent or in combination with anti-PD-1/PD-L1 antibodies has demonstrated tolerable safety profiles, favorable PK/PD profiles, and encouraging preliminary efficacy results.

